Heavy Neutral Lepton Searches at ICARUS-NuMI

JOSU HERNANDEZ-GARCIA

Based on arXiv: 2408.03383. In collaboration with A. De Roeck and A. Chatterjee

LIGHT DARK WORLD 2025 | SEPT 2025









Project CNS2022-136013 funded by MICIU/AEI/10.13039/501100011033 and the European Union NextGenerationEU/PRTR 2025 | Sept 2025










LLPs: Portals to BSM Physics


Long-Lived Particles (LLPs) could be the portal to hidden sectors addressing different Standard Model open problems





LLPs: Portals to BSM Physics


Long-Lived Particles (LLPs) could be the portal to hidden sectors addressing different Standard Model open problems



BSM portal	New particle	New parameters	SM open problem
Fermion portal	HNLs – N	$M_N, U_{lpha N}$	Neutrino masses
Axion portal	ALPs – a	m_a, f_a	Strong CP
Scalar portal	Dark scalar – S	$m_S, heta$	Vacuum stability
Vector portal	Dark photon – Z'	$m_{Z'}$, ϵ	DM candidate

LLPs: Portals to BSM Physics

Long-Lived Particles (LLPs) could be the portal to hidden sectors addressing different Standard Model open problems



BSM portal	New particle	New parameters	SM open problem
Fermion portal	HNLs – N	$M_N, U_{lpha N}$	Neutrino masses
Axion portal	ALPs – a	m_a, f_a	Strong CP
Scalar portal	Dark scalar – S	$m_S, heta$	Vacuum stability
Vector portal	Dark photon – Z'	$m_{Z'}, \epsilon$	DM candidate

LLPs signals at neutrino experiments

Neutrinos are feebly-interacting particles

- intense sources needed
- large detectors needed

Interactions \rightarrow scale with detector mass

LLPs decays → scale with detector volume

Generally lower backgrounds if

- Detector has low density/mass
- decays can be fully reconstructed ⇒
 instrumented detector (ECAL, magnetic field, ...)

Phenomenology of Heavy Neutral Leptons

Origin of neutrino masses and mixings

Minimal model: 2 HNLs needed. Phenomenology given by

4 free parameters:

- HNL mass, M_N
- mixing with three ν_{α} , $U_{\alpha N}$

HNL mixes with active neutrinos

$$\nu_{\alpha} = \sum_{i} U_{\alpha i} \nu_{i} + U_{\alpha N} N$$

Heavy Neutral Leptons (HNLs)

HNL CC and NC interactions:

$$\mathcal{L}_{N,int} \supset -\frac{m_W}{v} \overline{N} U_{\alpha N}^* \gamma^{\mu} \mathcal{L}_{L\alpha} W_{\mu}^+$$


$$-\frac{m_Z}{\sqrt{2} v} \overline{N} U_{\alpha N}^* \gamma^{\mu} \nu_{L\alpha} Z_{\mu}$$

$$U_{\alpha N} \ll 1 \Rightarrow$$
 weaker than weak interaction

Experimental results shown single flavor dominance \Rightarrow HNL coupled to one flavor at a time:

$$|U_{eN}|^2 : |U_{\mu N}|^2 : |U_{\tau N}|^2 = 1 : 0 : 0$$
 $|U_{eN}|^2 : |U_{\mu N}|^2 : |U_{\tau N}|^2 = 0 : 1 : 0$ $|U_{eN}|^2 : |U_{\mu N}|^2 : |U_{\tau N}|^2 = 0 : 0 : 1$ e dominance μ dominance τ dominance

Heavy Neutral Leptons: present bounds




Present bounds on the HNL mixing from ν oscillations, kinks in beta-decay, peaks in meson decays, beam-dumps, colliders, and non-unitarity of the PMNS.

https://github.com/mhostert/Heavy-Neutrino-Limits E. Fernandez-Martinez, M. Gonzalez-Lopez, JHG, M. Hostert, J. Lopez-Pavon. JHEP **09** 2023 001



04/20

Heavy Neutral Leptons: present bounds




Present bounds in the HNL mass window up to ~ 2 GeV led by peak searches and beam-dump experiments

https://github.com/mhostert/Heavy-Neutrino-Limits E. Fernandez-Martinez, M. Gonzalez-Lopez, JHG, M. Hostert, J. Lopez-Pavon. JHEP **09** 2023 001



Heavy Neutral Leptons: present bounds



Bounds from on-going and near future experiments.

The ICARUS experiment at SBN

Proposed to:

- Clarify LSND and MiniBooNE anomalies
- Explore sterile neutrino in the eV mass scale

Three liquid Ar TPCs

Taking data since 2015


SBND

270 tons
Baseline: 110 m

2024

2021

 μ BooNE



170 tons

Baseline: 470 m


2015 - 2021

Short-Baseline Neutrino (SBN)


ICARUS
760 tons
Baseline: 600 m

Exposed to the Booster Neutrino Beam (BNB) produced from 8 GeV protons from the booster

Experimental setup: SBN



Experimental setup: BNB and NuMI




ICARUS is also exposed to the NuMI ν -beam.


NuMI: 120 GeV protons from main injector against graphite target, accumulating $6.6 \cdot 10^{20}$ PoT/year.

C. Rubbia (ICARUS-WA104, LAr1-ND, MicroBooNE), (2015)


Experimental setup: BNB and NuMI



HNLs could be produced via meson decay at the beam-dump experiment beam




HNLs could be produced via meson decay at the beam-dump experiment beam



The number of decay events

$$N_{ev} = N_M \text{BR}(M \to N) \text{BR}(N \to \text{vis}) \epsilon_{det} \int dS \int dE_{\Psi} P(c\tau_N/M_N, E_N, \Omega_N) \frac{dn^{M \to N}}{dS dE_N}$$

HNLs could be produced via meson decay at the beam-dump experiment beam




The number of decay events

$$N_{ev} = N_M \text{BR}(M \to N) \text{BR}(N \to \text{vis}) \epsilon_{det} \int dS \int dE_{\Psi} P(c\tau_N/M_N, E_N, \Omega_N) \frac{dn^{M \to N}}{dS dE_N}$$

experiment-dependent quantities

HNLs could be produced via meson decay at the beam-dump experiment beam




The number of decay events


$$N_{ev} = N_M \text{BR}(M \to N) \text{BR}(N \to \text{vis}) \epsilon_{det} \int dS \int dE_{\Psi} P(c\tau_N/M_N, E_N, \Omega_N) \frac{dn^{M \to N}}{dS dE_N}$$

$$\text{HNL} \\ \text{production} \\ \text{HNL} \\ \text{decay} \\ \text{quantities}$$

HNLs could be produced via meson decay at the beam-dump experiment beam




HNLs could be produced via meson decay at the beam-dump experiment beam



Three sets of simulations are needed in order to estimate the number of events:

HNLs could be produced via meson decay at the beam-dump experiment beam




Three sets of simulations are needed in order to estimate the number of events:

- Meson production at the target station: we consider the mesons with relevant branching fraction to neutrinos
 - π^{\pm} & K^{\pm} : GEANT4 simulations including the interaction of 120 GeV protons on the graphite target, the production of mesons, and the re-interaction and focusing in the magnetic horns (G4NuMI)

L. Aliaga Soplin, Neutrino Flux Prediction for the NuMI Beamline, Ph.D. thesis, William-Mary Coll. (2016)

HNLs could be produced via meson decay at the beam-dump experiment beam




Three sets of simulations are needed in order to estimate the number of events:

- Meson production at the target station: we consider the mesons with relevant branching fraction to neutrinos
 - π^{\pm} & K^{\pm} : GEANT4 simulations including the interaction of 120 GeV protons on the graphite target, the production of mesons, and the re-interaction and focusing in the magnetic horns (G4NuMI)
 - D^{\pm} , D_s^{\pm} & τ^{\pm} : simulated with GEANT4 + Pythia8

P. Coloma, E. Fernandez-Martinez, M. Gonzalez-Lopez, J. Hernandez-Garcia, Z. Pavlovic, Eur. Phys. J. C 81 (2021) 1, 78

HNLs could be produced via meson decay at the beam-dump experiment beam




Three sets of simulations are needed in order to estimate the number of events:

- Meson production at the target station: we consider the mesons with relevant branching fraction to neutrinos
- MonteCarlo simulations of HNL production and HNL decay:


We use a FeynRules model file that contains effective op. describing interactions between light mesons and HNL. Interfacing the model file with an event generator allows for the simulation of fully differential event distributions.

The simulations of HNL fluxes were performed with HNLux:



P. Coloma, E. Fernandez-Martinez, M. Gonzalez-Lopez, J. Hernandez-Garcia, Z. Pavlovic, Eur. Phys. J. C 81 (2021) 1, 78


The HNL sensitivity line shows a characteristic shape coming from:



- Meson production yield N_M
 - proton *E*
 - luminosity (PoT)
 - target material/geometry

 $N_{\pi} > N_{K} > N_{D} \Rightarrow$ different regions

The HNL sensitivity line shows a characteristic shape coming from:




- Meson production yield N_M
 - proton *E*
 - luminosity (PoT)

 $N_{\pi} > N_{K} > N_{D} \Rightarrow$ different regions

- target material/geometry
- BR $(M \to N\ell_{\alpha}X) \Rightarrow M_N \leq M_M m_{\ell_{\alpha}} m_X$: upper limit of M_N

The HNL sensitivity line shows a characteristic shape coming from:




- Meson production yield N_M
 - proton E
 - luminosity (PoT)

different regions

 $N_{\pi} > N_{K} > N_{D} \Rightarrow$

- target material/geometry
- BR $(M \to N\ell_{\alpha}X) \Rightarrow M_N \leq M_M m_{\ell_{\alpha}} m_X$: upper limit of M_N
- BR($N \to \text{vis}$) $\Rightarrow M_N \ge \sum m_{\text{vis}}$: lower limit of M_N

The HNL sensitivity line shows a characteristic shape coming from:



- Meson production yield N_M
 - proton *E*
 - luminosity (PoT)
 - target material/geometry

$$N_{\pi} > N_{K} > N_{D} \Rightarrow$$

different regions


- BR $(M \to N\ell_{\alpha}X) \Rightarrow M_N \leq M_M m_{\ell_{\alpha}} m_X$: upper limit of M_N
- BR($N \to \text{vis}$) $\Rightarrow M_N \ge \sum m_{\text{vis}}$: lower limit of M_N

 $|U_{\tau N}|^2$ only via $D/D_s \to \tau$, and τ decays \Rightarrow less sensitivity due to suppressed production yield

HNL sensitivity: signal only

Following the Feldman and Cousins prescription for a Poisson distribution with no background:

$$N_{ev}^{ex} < 2.3 \Rightarrow 90\% \text{ CL}$$




- $1.32 \cdot 10^{21}$ PoT from NuMI beam $\Rightarrow \sim$ by end of 2026
- 100 % signal efficiency

HNL sensitivity: signal only

Following the Feldman and Cousins prescription for a Poisson distribution with no background:

$$N_{ev}^{ex} < 2.3 \Rightarrow 90\% \text{ CL}$$




- $1.32 \cdot 10^{21}$ PoT from NuMI beam $\Rightarrow \sim$ by end of 2026
- 100 % signal efficiency

HNL sensitivity: signal only

Following the Feldman and Cousins prescription for a Poisson distribution with no background:

$$N_{ev}^{ex} < 2.3 \Rightarrow 90\% \text{ CL}$$



- $1.32 \cdot 10^{21}$ PoT from NuMI beam $\Rightarrow \sim$ by end of 2026
- 100 % signal efficiency


The dominant background comes from the CC and NC SM neutrino interactions with the 40 Ar.

If we focus on HNL decay channels with higher BR, we can distinguish among the following three topologies:

The dominant background comes from the CC and NC SM neutrino interactions with the $^{40}\mathrm{Ar}$.

If we focus on HNL decay channels with higher BR, we can distinguish among the following three topologies:

• Channels with two μ -like tracks



Decay channel

$$N \to \pi^{+}\mu^{-}$$

$$N \to \rho^{+}\mu^{-}, \pi^{+}\pi^{0}\mu^{-}$$

$$N \to \rho^{0}\nu, \pi^{+}\pi^{-}\nu$$

 $N \to \mu^+ \mu^- \nu$

Main source of background


Single pion production from ν_{μ} resonant or coherent nucleus scattering

Charm production in CC deep inelastic ν_{μ} nucleus scattering (suppressed)

The dominant background comes from the CC and NC SM neutrino interactions with the 40 Ar.

If we focus on HNL decay channels with higher BR, we can distinguish among the following three topologies:

• Channels with one e-like track/shower and one μ -like track



Decay channel

 $N \to \pi^+ e^ N \to \rho^+ e^-, \pi^+ \pi^0 e^{-*}$

Main source of background

Single pion production from ν_e resonant or coherent nucleus scattering


 $N \rightarrow \mu^+ e^- \nu$

Charm production in CC deep inelastic ν_e or ν_μ nucleus scattering (suppressed)

The dominant background comes from the CC and NC SM neutrino interactions with the $^{40}\mathrm{Ar}$.


If we focus on HNL decay channels with higher BR, we can distinguish among the following three topologies:

• Channels with two *e*-like tracks/showers



Decay channel	Main source of background
$N \to \pi^0 \nu, 2\gamma \nu$	Single pion production from NC resonant nucleus scattering
$N \rightarrow e^+ e^- \nu$	Charm production in CC deep inelastic ν_e nucleus scattering (suppressed)

ν_{α} -⁴⁰Ar number of interactions



Number of ν_{α} -40 Ar interactions in the detector given by

$$n_{ev} = n_t \int \phi_{\nu_{\alpha}} \sigma_{\nu_{\alpha}}^{40Ar}$$
number of targets ICARUS

In one year:

	CC $ u_{lpha}$ - 40 Ar	NC ν_{α} -40Ar
$\alpha = e$	19142	2109
$\alpha = \mu$	341257	42610

Kinematical cuts

Momentum distributions of signal and background very different.

HNLs come from meson decays produced at the target \Rightarrow HNLs highly boosted:


- Low reconstructed transverse momentum p_T
- Small angle θ between μ -like tracks

Kinematical cuts

Momentum distributions of signal and background very different.

HNLs come from meson decays produced at the target \Rightarrow HNLs highly boosted:

- Low reconstructed transverse momentum p_T
- Small angle θ between μ -like tracks




By applying kinematical cuts:

- Bkg. rej. eff. up to $\simeq 90\%$
- Signal eff. up to $\simeq 92\%$

HNL sensitivity: signal + background


Combining the channels with two μ -like tracks



• Solid purple line: F&C analysis assuming no background

HNL sensitivity: signal + background


Combining the channels with two μ -like tracks



- Solid purple line: F&C analysis assuming no background
- Dashed purple line: unbinned Gaussian χ^2 analysis including both signal and bkg after event kinematical cuts

HNL sensitivity: signal + background


Combining the channels with two μ -like tracks



- Solid purple line: F&C analysis assuming no background
- Dashed purple line: unbinned Gaussian χ^2 analysis including both signal and bkg after event kinematical cuts
- Pink band: (15, 30) % signal efficiency cut to the negligible background case that covers the signal + bkg result

HNL sensitivity: ICARUS over present constraints

Combining leading HNL decay channels and following the Feldman and Cousins prescription \Rightarrow 90 % CL ICARUS-like expected bounds assuming 1.32 \cdot 10²¹ PoT




• Green line: signal only

• Pink band: (15, 30) % signal efficiency cut \Rightarrow conservative limit

HNL sensitivity: ICARUS over present constraints

Combining leading HNL decay channels and following the Feldman and Cousins prescription \Rightarrow 90 % CL ICARUS-like expected bounds assuming 1.32 \cdot 10²¹ PoT



• Pink band: (15, 30) % signal efficiency cut \Rightarrow conservative limit

Sensitivity results compatible with arXiv: 2409.04394

Summary

- We have computed, for the first time, the expected sensitivity of ICARUS to HNLs from NuMI.
- ICARUS shows potential for testing unexplored regions of the HNL parameter space.
- The ICARUS collaboration should analyze the data accumulated by the end of 2026 and perform a full dedicated analysis

Summary

- We have computed, for the first time, the expected sensitivity of ICARUS to HNLs from NuMI.
- ICARUS shows potential for testing unexplored regions of the HNL parameter space.
- The ICARUS collaboration should analyze the data accumulated by the end of 2026 and perform a full dedicated analysis











Project CNS2022-136013 funded by MICIU/AEI/10.13039/501100011033 and the European Union NextGenerationEU/PRTR 2025 | Sept 2025

This research is also supported by the EU H2020 research and innovation programme under the MSC grant agreement No 860881-11 HIDDeN, as well as by the Spanish Ministerio de Ciencia e Innovación project PID2020-113644GB-loo, the Spanish Research Agency (Agencia Estatal de Investigación) through the project CNS2022-136013, and Severo Ochoa Excellence Program CEX2023-001292-S.

Back-up

Status of NuMI after 2024 power failure

A major power failure of a transformer on the NuMI beam line at the end of August 2024 stopped its activity.

As a result, NuMI is not running, and its continuation until the end of 2026 is not yet clear.


From 2027 on, Fermilab will focus on the LBNF beam line for DUNE.

ICARUS has so far collected $\sim 6 \cdot 10^{20} \, \text{PoT} \Rightarrow \text{roughly half of what was assumed for our results } (1.32 \cdot 10^{21} \, \text{PoT}).$

If NuMI resumes its activity during 2026, ICARUS could still collect $\sim 1.32 \cdot 10^{21}$ PoT.

Otherwise, our sensitivity results should be corrected by a factor of $\sim \sqrt{2}$.

Heavy Neutral Leptons: present bounds




Present bounds in the HNL mixing from ν oscillations, kinks in beta-decay, peaks in meson decays, beam-dumps, colliders, and non-unitarity of the PMNS.

https://github.com/mhostert/Heavy-Neutrino-Limits E. Fernandez-Martinez, M. Gonzalez-Lopez, JHG, M. Hostert, J. Lopez-Pavon. JHEP **09** 2023 001

LDW2025



Heavy Neutral Leptons: present bounds




Present bounds in the HNL mixing from ν oscillations, kinks in beta-decay, peaks in meson decays, beam-dumps, colliders, and non-unitarity of the PMNS.

https://github.com/mhostert/Heavy-Neutrino-Limits E. Fernandez-Martinez, M. Gonzalez-Lopez, JHG, M. Hostert, J. Lopez-Pavon. JHEP **09** 2023 001



Heavy Neutral Leptons: present bounds



Present bounds in the HNL mixing from ν oscillations, kinks in beta-decay, peaks in meson decays, beam-dumps, colliders, and non-unitarity of the PMNS.

https://github.com/mhostert/Heavy-Neutrino-Limits E. Fernandez-Martinez, M. Gonzalez-Lopez, JHG, M. Hostert, J. Lopez-Pavon. JHEP **09** 2023 001



Heavy Neutral Leptons: production and decay

HNL production

Leptonic and semileptonic meson decays

	$\pi \rightarrow$	$K^0 \rightarrow$	$K \rightarrow$	$D \rightarrow$	$D_s \rightarrow$	au ightarrow
$ U_{eN} ^2$	eN		eN	eN	eN	
	_	$\pi^0 e N$	$\pi e N$	K^0eN		
$ U_{\mu N} ^2$	μN		$\overline{\mu N}$	μN	$\overline{\mu N}$	
		$\pi^0 \mu N$	$\pi \mu N$	$K^0\mu N$		
$ U_{ au N} ^2$	_			au N	au N	πN
	_					$\pi\pi^0 N$
						e u N
	_					$\mu u N$

	π	K	D	D_s	au
P^+/PoT	6.0	1.1	$1.2\cdot 10^{-5}$	$3.3\cdot 10^{-6}$	$2.1\cdot 10^{-7}$
P^-/PoT	4.1	0.4	$1.9\cdot 10^{-5}$	$4.6\cdot 10^{-6}$	$3.0\cdot 10^{-7}$

Table I. Averaged parent production yield at the NuMI target per PoT during its neutrino mode. A pre-selection of light mesons that would produce neutrinos towards the region where ICARUS is placed has been done.

• HNL decay
Visible HNL decay channels with higher branching ratio

			$N \rightarrow$		
$ U_{eN} ^2$	$ee\nu$	$\mu\mu\nu$	$e\mu\nu$	e hadr.	ν hadr.
	$ u \pi^0 $	$e\pi$	$e\rho$		
$ II _{N} ^{2}$	$ee\nu$	$\mu\mu\nu$	$e\mu\nu$	μ hadr.	ν hadr.
$\mu \nu$	$ u \pi^0 $	$\mu\pi$	μho		
$ U_{\tau N} ^2$	$ee\nu$	$\mu\mu\nu$			ν hadr.
$ \cup \tau N $	$ u \pi^0 $			$ u ho^0$	

Statistical analisys

Feldman and Cousins

Prescription for a Poisson distribution with no background and under the hypothesis of no events being observed

$$N^{ex}(\phi) < 2.44$$
, for 90 % CL


• Unbinned Gaussian χ^2

$$\chi^{2} = min_{\xi} \left\{ \left(\frac{N^{ex}(\phi, \xi) - N^{ob}(\xi)}{\sigma_{N}(\xi)} \right)^{2} + \left(\frac{\xi}{\sigma_{f}} \right)^{2} \right\}$$

where $N^{ex}(\phi,\xi)$ is the total expected event rate including both signal and background events, and $N^{ob}(\xi)$ is the observed events; with the uncertainty on the flux normalization, σ_f , taken as $20\,\%$, and $\sigma_N(\xi) = \sqrt{N^{ob}(\xi)}$ is the statistical error.


$$\chi^2 < 4.61$$
, for 90 % CL

Kinematical cuts




Present and nHNL sensitivity: present and near-future

Expected sensitivities assuming signal only (no background)



2 years of ICARUS NuMI (2026). 4.5 years of DUNE Phase II (2040). 15 years of SHiP (2045).

The SHiP experiment




SHiP experiment highlights:

- Proton energy of 400 GeV
- Expected running time $\simeq 2030 2045$
- Large beam intensity $4 \cdot 10^{19}$ PoT/year \Rightarrow an expected total of $\simeq 6 \cdot 10^{20}$ PoT
- "Background-free" experiment for many scenarios
- Fully instrumented

designed for LLP searches


HNLs searches: DUNE ND-GAr vs SHiP



Better sensitivity of DUNE ND-GAr in the π and K regions

Maksym Ovchynnikov - Search for new physics at SHiP (IFIC seminar)

HNLs searches: DUNE ND-GAr vs SHiP




Better sensitivity of DUNE ND-GAr in the π and K regions

Better sensitivity of SHiP in the D and B regions

Maksym Ovchynnikov - Search for new physics at SHiP (IFIC seminar)

HNLs searches: DUNE ND-GAr vs SHiP



Better sensitivity of DUNE ND-GAr in the π and K regions Better sensitivity of SHiP in the D and B regions Similar results for the $|U_{\mu N}|^2$. Better sensitivity of SHiP in the $|U_{\tau N}|^2$ since it comes from D decays.

Maksym Ovchynnikov - Search for new physics at SHiP (IFIC seminar)